

# t9\_flang\_3 (TMEhikdR- FuVu14888fU8hNCLMwTeJWRZSot)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k8\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_flang\_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_catalan2 : \iota \Rightarrow \iota$  be given. Let  $k7\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow ((X1 \in k8\_flang\_1 X0 X2) \Leftrightarrow (\exists X3. (v7\_ordinal1 X3) \wedge (X1 \in k7\_flang\_1 X0 X2 X3))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow (\forall X3. (v7\_ordinal1 X3) \Rightarrow ((X1 \in k1\_flang\_3 X0 X2 X3) \Leftrightarrow (\exists X4. (v7\_ordinal1 X4) \wedge ((r1\_xxreal\_0 X3 X4) \wedge (X1 \in k7\_flang\_1 X0 X2 X4)))))) \quad (2)$$

Assume the following.

$$\forall X0. k3\_catalan2 X0 = k8\_afinsq\_1 X0 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (4)$$

## Theorem 1

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow (\forall X2. (v7\_ordinal1 X2) \Rightarrow (r1\_tarski (k1\_flang\_3 X0 X1 X2) (k8\_flang\_1 X0 X1)))$$