

## t9\_funct\_2

(TMHJN3q4x1cFm7gHJmtmq5LKLFMGtxrqpb1)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\ & X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))))) \Rightarrow \\ & (((X1 = k1\_xboole\_0) \Rightarrow (X0 = k1\_xboole\_0)) \Rightarrow (X2 \in k1\_funct\_2 X0 X1)) \end{aligned} \tag{1}$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 X0 X0) \wedge \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X0)))))) \Rightarrow (X1 \in k1\_funct\_2 \\ & X0 X0) \end{aligned}$$