

t9\_funct\_3 (TM-  
LYda4mT9smCanyGyeMZy9C2oaA56JJ6At)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_3 : \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((X0 \in k9\_xtuple\_0 (k1\_funct\_3 X1)) \Rightarrow (k1\_funct\_1 (k1\_funct\_3 X1) X0 = k7\_relat\_1 X1 X0)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 X1) \Rightarrow ((v1\_xboole\_0 X1) \vee (X0 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (v1\_relat\_1 X1) \Rightarrow (r1\_tarski (k7\_relat\_1 X1 X0) (k10\_xtuple\_0 X1)) \quad (4)$$

Assume the following.

$$\forall X0. \neg v1\_xboole\_0 (k1\_zfmisc\_1 X0) \quad (5)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow ((v1\_relat\_1 (k1\_funct\_3 X0)) \wedge (v1\_funct\_1 (k1\_funct\_3 X0))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(r1\_tarSKI X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow (X2 \in X1)) \quad (7)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge(v1\_funct\_1 X0))\Rightarrow(\forall X1.(X1 = k10\_xtuple\_0 X0)\Leftrightarrow(\forall X2.(X2 \in X1)\Leftrightarrow(\exists X3.(X3 \in k9\_xtuple\_0 X0)\wedge(X2 = k1\_funct\_1 X0 X3)))) \quad (8)$$

**Theorem 1**

$$\forall X0.((v1\_relat\_1 X0)\wedge(v1\_funct\_1 X0))\Rightarrow(r1\_tarSKI (k10\_xtuple\_0 (k1\_funct\_3 X0)) (k1\_zfmisc\_1 (k10\_xtuple\_0 X0)))$$