

## t9\_funct\_9

(TMXqgh7SnRaxTokRmvfpaisqAMWA26MQAxk)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v3\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_9 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k13\_valued\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_valued\_0 X0))) \Rightarrow \\ & (\forall X1.(v1\_xcmplx\_0 X1) \Rightarrow ((k9\_xtuple\_0 (k13\_valued\_1 X0 \\ & X1) = k9\_xtuple\_0 X0) \wedge (\forall X2.(X2 \in k9\_xtuple\_0 X0) \Rightarrow (k1\_funct\_1 \\ & (k13\_valued\_1 X0 X1) X2 = k6\_xcmplx\_0 (k1\_funct\_1 X0 X2) X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_valued\_0 \\ & X0))) \wedge (v1\_xcmplx\_0 X1)) \Rightarrow ((v1\_relat\_1 (k13\_valued\_1 X0 X1)) \wedge \\ & (v1\_funct\_1 (k13\_valued\_1 X0 X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 \\ & X1)) \Rightarrow ((v1\_funct\_9 X1 X0) \Leftrightarrow ((X0 \neq k6\_numbers) \wedge (\forall X2.(v1\_xreal\_0 \\ & X2) \Rightarrow (((X2 \in k9\_xtuple\_0 X1) \Rightarrow (k2\_xcmplx\_0 X2 X0 \in k9\_xtuple\_0 X1)) \wedge \\ & (((k2\_xcmplx\_0 X2 X0 \in k9\_xtuple\_0 X1) \Rightarrow (X2 \in k9\_xtuple\_0 X1)) \wedge \\ & (X2 \in k9\_xtuple\_0 X1) \Rightarrow (k1\_funct\_1 X1 X2 = k1\_funct\_1 X1 (k2\_xcmplx\_0 \\ & X2 X0)))))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v3\_valued\_0 X0)) \Rightarrow ((v1\_relat\_1 X0) \wedge (v1\_valued\_0 X0)) \quad (4)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (5)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & ((v1\_relat\_1 X2) \wedge ((v1\_funct\_1 X2) \wedge (v3\_valued\_0 X2))) \Rightarrow ((v1\_funct\_9 \\ & X2 X0) \Rightarrow (v1\_funct\_9 (k13\_valued\_1 X2 X1) X0)))) \end{aligned}$$