

t9_glib_000 (TMaNKVonjML- BkfkV4ArPvUArimFk942TH9)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k13_glib_000 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (X0 \in k9_xtuple_0 (k16_funcop_1 X1 X2)) \Rightarrow (X0 = X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((\neg X0 \in k9_xtuple_0 X1) \Rightarrow (k1_funct_1 (k1_funct_4 X2 X1) X0 = k1_funct_1 X2 X0))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k16_funcop_1 X0 X1)) \wedge (v1_funct_1 (k16_funcop_1 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0)))) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow (\forall X2. k13_glib_000 X0 X1 X2 = k1_funct_4 X0 (k16_funcop_1 X1 X2))) \quad (4)$$

Theorem 1

$$\forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0)))) \Rightarrow (\forall X1. \forall X2. (v7_ordinal1 X2) \Rightarrow (\forall X3. (v7_ordinal1 X3) \Rightarrow ((X2 \neq X3) \Rightarrow (k1_funct_1 X0 X3 = k1_funct_1 (k13_glib_000 X0 X2 X1) X3))))$$