

t9_nat_5 (TMTCHt- nYHy1FqL9k1ykpykrW9rLyYWAJck1)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_rvsum_1 : \iota \Rightarrow \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (\forall X1. (v1_relat_1 X1) \Rightarrow ((r1_tarski (k10_xtuple_0 X0) (k9_xtuple_0 X1)) \Rightarrow (k9_xtuple_0 (k3_relat_1 X0 X1) = k9_xtuple_0 X0))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v5_relat_1 X1 X0)) \Rightarrow (k2_relset_1 X0 X1 = k10_xtuple_0 X1) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \Rightarrow (k1_rvsum_1 X0 = k10_xtuple_0 X0) \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. \forall X5. \\ & (((v1_funct_1 X4) \wedge (m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))) \wedge ((v1_funct_1 X5) \wedge (m1_subset_1 X5 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X2 X3)))))) \Rightarrow (k1_partfun1 X0 X1 X2 X3 X4 X5 = k3_relat_1 X4 X5) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0. \forall X1. v1_relat_1 (k2_zfmisc_1 X0 X1) \quad (8)$$

Assume the following.

$$v3_membered k1_numbers \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge ((v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (10)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \quad (11)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (12)$$

Assume the following.

$$\forall X0. \forall X1. v1_relat_1 (k3_relat_1 X0 X1) \quad (13)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v5_relat_1 X1 X0)) \Rightarrow (m1_subset_1 (k2_relset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. \forall X5. \\ & (((v1_funct_1 X4) \wedge (m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))) \wedge ((v1_funct_1 X5) \wedge (m1_subset_1 X5 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X2 X3)))))) \Rightarrow ((v1_funct_1 (k1_partfun1 X0 X1 X2 X3 X4 X5)) \wedge (m1_subset_1 \\ & (k1_partfun1 X0 X1 X2 X3 X4 X5) (k1_zfmisc_1 (k2_zfmisc_1 X0 X3)))) \end{aligned} \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1)))\Rightarrow((m1_finseq_1 X1 X0)\Leftrightarrow(r1_tarski (k10_xtuple_0 X1) X0)) \quad (16)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0)\Rightarrow((v1_finseq_1 X0)\Leftrightarrow(\exists X1.(v7_ordinal1 X1)\wedge(k9_xtuple_0 X0 = k2_finseq_1 X1))) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (18)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_relat_1 X1)) \quad (19)$$

Assume the following.

$$\forall X0.(v3_membered X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v3_membered X1)) \quad (20)$$

Assume the following.

$$\forall X0.\forall X1.(v3_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v3_valued_0 X2)) \quad (21)$$

Theorem 1

$$\forall X0.\forall X1.(m2_finseq_1 X1 k5_numbers)\Rightarrow(\forall X2.(m2_finseq_1 X2 X0)\Rightarrow((r1_tarski (k1_rvsum_1 X1) (k4_finseq_1 X2))\Rightarrow(m2_finseq_1 (k1_partfun1 k5_numbers k5_numbers k5_numbers X0 X1 X2) X0)))$$