

t9_pzfmisc1 (TMNMRoDihCMP- PRKVqmzaEW2gUhMDiXuvdfD)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r6_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_pzfmisc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
& \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 X0) \wedge \\
& (v1_funct_1 X1) \wedge (v1_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1_relat_1 \\
& X2) \wedge ((v4_relat_1 X2 X0) \wedge ((v1_funct_1 X2) \wedge (v1_partfun1 X2 X0)))) \Rightarrow \\
& (\forall X3. ((v1_relat_1 X3) \wedge ((v4_relat_1 X3 X0) \wedge ((v1_funct_1 \\
& X3) \wedge (v1_partfun1 X3 X0)))) \Rightarrow ((r6_pboole X0 X1 (k2_pzfmisc1 X0 X2 \\
& X3)) \Rightarrow (\forall X4. ((v1_relat_1 X4) \wedge ((v4_relat_1 X4 X0) \wedge ((v1_funct_1 \\
& X4) \wedge (v1_partfun1 X4 X0)))) \Rightarrow (((r6_pboole X0 X4 X2) \vee (r6_pboole \\
& X0 X4 X3)) \Rightarrow (r1_pboole X0 X4 X1))))))
\end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0. \forall X1. \forall X2. (((v1_relat_1 X1) \wedge ((v4_relat_1 \\
& X1 X0) \wedge ((v1_funct_1 X1) \wedge (v1_partfun1 X1 X0)))) \wedge ((v1_relat_1 \\
& X2) \wedge ((v4_relat_1 X2 X0) \wedge ((v1_funct_1 X2) \wedge (v1_partfun1 X2 X0)))) \Rightarrow \\
& (r6_pboole X0 X1 X1)
\end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0. \forall X1. \forall X2. (((v1_relat_1 X1) \wedge ((v4_relat_1 \\
& X1 X0) \wedge ((v1_funct_1 X1) \wedge (v1_partfun1 X1 X0)))) \wedge ((v1_relat_1 \\
& X2) \wedge ((v4_relat_1 X2 X0) \wedge ((v1_funct_1 X2) \wedge (v1_partfun1 X2 X0)))) \Rightarrow \\
& ((r6_pboole X0 X1 X2) \Leftrightarrow (X1 = X2))
\end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.(((v1_relat_1 X1)\wedge((v4_relat_1 \\
& X1 X0)\wedge((v1_funct_1 X1)\wedge(v1_partfun1 X1 X0))))\wedge((v1_relat_1 \\
& X2)\wedge((v4_relat_1 X2 X0)\wedge((v1_funct_1 X2)\wedge(v1_partfun1 X2 X0))))\Rightarrow \\
& ((v1_relat_1 (k2_pzfmisc1 X0 X1 X2))\wedge((v4_relat_1 (k2_pzfmisc1 \\
& X0 X1 X2) X0)\wedge((v1_funct_1 (k2_pzfmisc1 X0 X1 X2))\wedge(v1_partfun1 \\
& (k2_pzfmisc1 X0 X1 X2) X0))))
\end{aligned} \tag{4}$$

Theorem 1

$$\begin{aligned}
& \forall X0.\forall X1.((v1_relat_1 X1)\wedge((v4_relat_1 X1 X0)\wedge(\\
& (v1_funct_1 X1)\wedge(v1_partfun1 X1 X0))))\Rightarrow(\forall X2.((v1_relat_1 \\
& X2)\wedge((v4_relat_1 X2 X0)\wedge((v1_funct_1 X2)\wedge(v1_partfun1 X2 X0))))\Rightarrow \\
& (\forall X3.((v1_relat_1 X3)\wedge((v4_relat_1 X3 X0)\wedge((v1_funct_1 \\
& X3)\wedge(v1_partfun1 X3 X0))))\Rightarrow(((r6_pboole X0 X1 X2)\vee(r6_pboole \\
& X0 X1 X3))\Rightarrow(r1_pboole X0 X1 (k2_pzfmisc1 X0 X2 X3))))
\end{aligned}$$