

t9_scmyciel (TMXKQMwC-R- WHRf5BpJw1Z56sWjVwBkYBzH1q)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k1_tarSKI : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_zfmisc_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v3_card_1 X1 np_1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0))) \Rightarrow (\exists X2. (m1_subset_1 \\ X2 X0) \wedge (X1 = k1_tarSKI X2))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \Rightarrow ((m1_subset_1 X1 X0) \Leftrightarrow \\ (X1 \in X0))) \wedge ((v1_xboole_0 X0) \Rightarrow ((m1_subset_1 X1 X0) \Leftrightarrow (v1_xboole_0 \\ X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. (v3_card_1 X0 np_1) \Rightarrow ((\neg v1_xboole_0 X0) \wedge (v1_zfmisc_1 X0)) \quad (3)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (4)$$

Theorem 1

$$\begin{aligned} \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\neg (v3_card_1 \\ X1 np_1) \wedge (\forall X2. \neg (X2 \in X0) \wedge (X1 = k1_tarSKI X2))) \end{aligned}$$