

t9_seqm_3

(TMJs2ZpvViAEarCx2J7qhTxna7BfANebCY3)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_valued_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ & ((v5_valued_0 X0) \Leftrightarrow (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\ & (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow (\neg(\neg r1_xxreal_0 \\ & X2 X1) \wedge (r1_xxreal_0 (k8_nat_1 k1_numbers X0 X2) (k8_nat_1 k1_numbers \\ & X0 X1)))))) \end{aligned} \tag{1}$$

Assume the following.

$$m2_subset_1 k6_numbers k1_numbers k5_numbers \tag{2}$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ & ((v5_valued_0 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\ & (\neg(\neg r1_xxreal_0 X1 k6_numbers) \wedge (r1_xxreal_0 (k8_nat_1 k1_numbers \\ & X0 X1) (k8_nat_1 k1_numbers X0 k6_numbers)))))) \end{aligned}$$