

## t9\_waybel\_3

(TMN3k1BEATNtMe7qV4xT5s8P8DcBUj5puFE)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v3\_orders\_2 : \iota \Rightarrow o$  be given. Let  $v5\_orders\_2 : \iota \Rightarrow o$  be given. Let  $l1\_orders\_2 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $r2\_lattice3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_waybel\_3 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_waybel\_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r3\_orders\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_orders\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v3\_orders\_2 X0) \wedge (l1\_orders\_2 \\ & X0))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (\forall X2. \\ & (m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow ((X1 \in k1\_waybel\_3 X0 X2) \Leftrightarrow (r1\_waybel\_3 \\ & X0 X1 X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v3\_orders\_2 X0) \wedge ((v5\_orders\_2 \\ & X0) \wedge (l1\_orders\_2 X0)))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 \\ & X0)) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow ((r1\_waybel\_3 \\ & X0 X1 X2) \Rightarrow (r3\_orders\_2 X0 X1 X2)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(((\neg v2\_struct\_0 X0) \wedge ((v3\_orders\_2 \\ & X0) \wedge (l1\_orders\_2 X0))) \wedge ((m1\_subset\_1 X1 (u1\_struct\_0 X0)) \wedge ( \\ & m1\_subset\_1 X2 (u1\_struct\_0 X0)))) \Rightarrow ((r3\_orders\_2 X0 X1 X2) \Leftrightarrow (r1\_orders\_2 \\ & X0 X1 X2)) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.(l1\_orders\_2 X0) \Rightarrow (\forall X1.\forall X2.(m1\_subset\_1 \\ & X2 (u1\_struct\_0 X0)) \Rightarrow ((r2\_lattice3 X0 X1 X2) \Leftrightarrow (\forall X3.(m1\_subset\_1 \\ & X3 (u1\_struct\_0 X0)) \Rightarrow ((X3 \in X1) \Rightarrow (r1\_orders\_2 X0 X3 X2)))) \end{aligned} \tag{4}$$

**Theorem 1**

$$\forall X0.((\neg v2\_struct\_0 X0) \wedge (v3\_orders\_2 X0) \wedge ((v5\_orders\_2 X0) \wedge (l1\_orders\_2 X0))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (r2\_lattice3 X0 (k1\_waybel\_3 X0 X1) X1))$$