

t9_yellow15 (TMFQfRrDmZRMMNiVmQd- pCwLnhdrb1qY6KKp)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_yellow15 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_setfam_1 : \iota \Rightarrow \iota$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k14_uncop_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_margrel1 : \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((X1 = k9_finseq_1 X0) \Leftrightarrow ((k3_finseq_1 X1 = np_1) \wedge (k10_xtuple_0 X1 = k1_tarski X0))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (2)$$

Assume the following.

$$\forall X0. k9_setfam_1 X0 = k1_zfmisc_1 X0 \quad (3)$$

Assume the following.

$$\forall X0. k9_finseq_1 X0 = k5_finseq_1 X0 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k12_finseq_1 X0 X1 = k5_finseq_1 X1) \quad (5)$$

Assume the following.

$$\forall X0. v1_finseq_1 (k5_finseq_1 X0) \quad (6)$$

Assume the following.

$$\forall X0. \neg v1_xboole_0 (k1_zfmisc_1 X0) \quad (7)$$

Assume the following.

$$\forall X0. (v1_relat_1 (k9_finseq_1 X0)) \wedge (v1_funct_1 (k9_finseq_1 X0)) \quad (8)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_finseq_1 X1 (k9_setfam_1 X0)) \wedge (m1_finseq_1 X2 k6_margrel1)) \Rightarrow (m2_finseq_1 (k1_yellow15 X0 X1 X2) (k9_setfam_1 X0)) \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (m2_finseq_1 (k12_finseq_1 X0 X1) X0) \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2_finseq_1 X1 (k9_setfam_1 X0)) \Rightarrow (\forall X2. \\ & (m2_finseq_1 X2 k6_margrel1) \Rightarrow (\forall X3. (m2_finseq_1 X3 (k9_setfam_1 X0)) \Rightarrow ((X3 = k1_yellow15 X0 X1 X2) \Leftrightarrow ((k3_finseq_1 X3 = k3_finseq_1 X1) \wedge (\forall X4. (v7_ordinal1 X4) \Rightarrow ((X4 \in k4_finseq_1 X1) \Rightarrow (k1_funct_1 X3 X4 = k14_funcop_1 (k1_funct_1 X2 X4) k8_margrel1 (k1_funct_1 X1 X4) (k6_subset_1 X0 (k1_funct_1 X1 X4)))))))))) \quad (11) \end{aligned}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. \\ & (m2_finseq_1 X2 k6_margrel1) \Rightarrow (k3_finseq_1 (k1_yellow15 X0 (k12_finseq_1 (k1_zfmisc_1 X0) X1) X2) = np_1)) \end{aligned}$$