

t9_zfmisc_1

(TMKAZk6XPUrfZAUStmGbV8715a5Qaz7Tjr)

October 27, 2020

Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (k2_xboole_0 X0 X1 = X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2_tarski X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 = X0) \vee (X3 = X1))) \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. k2_xboole_0 (k1_tarski X0) (k2_tarski X0 X1) = k2_tarski X0 X1$$